

# QUICK GUIDE FOR MANAGEMENT OF PATIENTS WITH COVID19: HOSPITALIST GUIDE + INITIAL WORK-UP

## FIRST STEPS: \*use interpreter phone if English is not first language\*

- At admission: HCP form +/- MOLST filled out and updated
- Attending to discuss realistic goals re. intubation and CPR
- Check baseline EKG

## LAB WORK-UP:

- Covid19 PCR testing (Biothreats p#30331) + Rapid viral panel

### At admission →

CBC with differential, BMP, Magnesium, CRP, LFT, CPK, LDH, PTT, INR procalcitonin, troponin, NT-proBNP, d-dimer

### Daily →

CBC with differential, BMP, Magnesium  
If patient is in ICU add: troponin, CPK

### Every other day →

LFT, CPK, troponin, CRP, LDH, d-dimer,  
If patient on propofol add: triglyceride

### If clinically worse →

LFT, CPK, troponin, CRP, procalcitonin, LDH, ferritin, d-dimer, fibrinogen, PTT, INR

## LAB RESULTS TO EXPECT: \*potential marker of disease severity

Normal WBC	Elevated AST*/ALT*
Lymphopenia*	Elevated CRP*
Mild thrombocytopenia	Elevated LDH*
BMP with elevated Cr	Elevated d-dimer*
Normal procalcitonin	Elevated troponin*

## RESPIRATORY CARE: See [Respiratory Failure Quick Guide](#) for details

@6L/min NC (goal SpO2 92 - 96% or PaO2 >75)

- Consult anesthesiology: contingency plan re. intubation
- Call Respiratory Therapy: consider venturi /nonrebreather
- Consult COVID ICU Triage (p39999): plan ICU transfer

**\*\*if respiratory decompensation or rapid increase in FiO2 call**

**anesthesiology to intubate: COVID airway pager 39265\*\***

Version 3.26.2020

To check for the most up to date recommendations, please visit the [full manual](#) or use the QR code here →

For urgent questions please consult the BWH ICU triage pager (#39999)

## ISOLATION: Remember these basics for covid + or rule-out patients

- Contact (gown + gloves) + Droplet (mask + eye protection)
- If aerosolizing procedure or ICU patient use N95 mask
- Aerosolizing procedures in negative pressure room only
- Avoid unnecessary aerosolizing procedures e.g. nebulization (switch to inhalers), high flow nasal canula, non-invasive ventilation (CPAP, BiPAP)
- OK to continue chronic night-time non-invasive ventilation, switch to BWH mask + machine because less aerosol risk

## CONSULTS to CALL: Upfront consults or when to call

- INFECTIOUS DISEASE → on ALL patients (discuss therapies)
- ANESTHESIOLOGY → if @6L/min NC or rapidly increasing FiO2
- RESPIRATORY THERAPY → if requiring 6L/min NC O2
- ICU TRIAGE → @6L/min NC or if concern for clinical worsening
- CARDIOLOGY → if concern for new heart failure, ACS, VT/VF, or cardiogenic shock
- ONCOLOGY → call primary oncologist at time of admission

## INITIAL MANAGEMENT CONSIDERATIONS:

**CT chest:** NOT necessary for diagnosis, recommend minimizing use of CT given challenges with isolation and transport

**Daily CXR:** NOT necessary unless it will change management plan

**IV fluids:** Conservative fluid management is important to mitigate risk of progression of respiratory failure

**Steroids:** Avoid using empirically, only use if other indication

**Antibiotics:** Follow BWH guidelines for empiric antibiotics based on patient risk factors, talk to ID consult about concerns

**Code Blue:** For covid + or covid rule-out, tell page operator this is covid patient; use normal protocol for donning of PPE prior to entering room, even if this delays CPR.



SCAN ME